

# 8 Analytic Trigonometry

- 8.1 Trigonometric Identities and Their Verification
- 8.2 The Addition and Subtraction Formulas
- 8.3 Double-Angle and Half-Angle Formulas
- 8.4 The Product–Sum Formulas
- 8.5 Trigonometric Equations



Much of the language and terminology of algebra carries over to trigonometry. For example, we have seen that algebraic expressions involve variables, constants, and algebraic operations. **Trigonometric expressions** involve these same elements but also permit trigonometric functions of variables and constants. They also allow algebraic operations upon these trigonometric functions. Thus,

$$x + 1 \sin x \quad \sin x + 1 \tan x \quad \frac{1 - \cos x}{\sec^2 x}$$

are all examples of trigonometric expressions.

The distinction between an identity and an equation also carries over to trigonometry. Thus, a **trigonometric identity** is true for all values that may be assumed by the variable, but a **trigonometric equation** is true only for certain values of the variable, called *solutions*. (Note that the solutions of a trigonometric equation may be expressed as real numbers or as angles.) As usual, the set of all solutions of a trigonometric equation is called the *solution set*.



## 8.1 Trigonometric Identities and Their Verification

### Fundamental Identities

In Section 7.3, we established the identity

$$\sin^2 t + \cos^2 t = 1 \quad (1)$$

If  $\cos t \neq 0$ , we may divide both sides of Equation (1) by  $\cos^2 t$  to obtain

$$\frac{\sin^2 t}{\cos^2 t} + \frac{\cos^2 t}{\cos^2 t} = \frac{1}{\cos^2 t}$$

or

$$\tan^2 t + 1 = \sec^2 t \quad (2)$$

Similarly, if  $\sin t \neq 0$ , dividing Equation (1) by  $\sin^2 t$  yields

$$\frac{\sin^2 t}{\sin^2 t} + \frac{\cos^2 t}{\sin^2 t} = \frac{1}{\sin^2 t}$$

or

$$\cot^2 t + 1 = \csc^2 t \quad (3)$$

Observe that  $\tan t$  and  $\sec t$  are undefined for exactly those values of  $t$  for which  $\cos t = 0$ . Similarly,  $\cot t$  and  $\csc t$  are undefined for those values of  $t$  for which  $\sin t = 0$ . It follows that Equations (2) and (3) are also identities. These identities, together with those discussed in Section 7.4, are called the **fundamental identities**. We summarize them in Table 1.

The fundamental identities can be used to simplify trigonometric expressions as well as to verify trigonometric identities. The purpose of including this topic is to present additional techniques that may be helpful in these situations. Such manipulations may enable us to see relationships that would otherwise be obscured.

The preferred method of verifying an identity is to transform one side of the equation into the other. Although we will use this method whenever practical, we recognize that it is also acceptable to transform each side independently, with the hope of arriving at the same expression and then showing that

the process can be reversed. This is often the technique used when both sides of the equation involve complicated expressions.

Unfortunately, we cannot present a specific set of steps that will “always work” to transform one side into the other. In fact, there are often many ways to handle a given identity. We will, however, demonstrate different techniques that we have found to be of value. We list the following suggestions on how to proceed, with examples to follow.

**TABLE 1** Fundamental Identities

$$\tan t = \frac{\sin t}{\cos t}$$

$$\cot t = \frac{\cos t}{\sin t}$$

$$\csc t = \frac{1}{\sin t}$$

$$\sec t = \frac{1}{\cos t}$$

$$\cot t = \frac{1}{\tan t}$$

$$\sin^2 t + \cos^2 t = 1$$

$$\tan^2 t + 1 = \sec^2 t$$

$$\cot^2 t + 1 = \csc^2 t$$

$$\sin t = \frac{1}{\csc t}$$

$$\cos t = \frac{1}{\sec t}$$

$$\tan t = \frac{1}{\cot t}$$

$$\sin^2 t = 1 - \cos^2 t$$

$$\cos^2 t = 1 - \sin^2 t$$

$$\tan^2 t = \sec^2 t - 1$$

$$\cot^2 t = \csc^2 t - 1$$

1. Factoring may help to simplify an expression.
2. It is often helpful to write all of the trigonometric functions in terms of sine and cosine.
3. Consider beginning with the more complicated expression and perform some of the indicated operations.
4. If you have the ratio of two trigonometric functions, it may be worthwhile to multiply both numerator and denominator by some trigonometric expression to obtain forms such as  $1 - \sin^2 \theta$ ,  $1 - \cos^2 \theta$ , or  $\sec^2 \theta - 1$ , which can be further simplified.

**EXAMPLE 1** USING TRIGONOMETRIC IDENTITIES

Simplify the expression  $\sin^2 x + \sin^2 x \tan^2 x$ .

**SOLUTION**

We begin by noting that  $\sin^2 x$  appears in both terms, which suggests that we factor.

$$\begin{aligned} \sin^2 x + \sin^2 x \tan^2 x &= \sin^2 x (1 + \tan^2 x) && \text{Factoring} \\ &= \sin^2 x \sec^2 x && 1 + \tan^2 x = \sec^2 x \\ &= \sin^2 x \left( \frac{1}{\cos^2 x} \right) && \sec x = \frac{1}{\cos x} \\ &= \tan^2 x && \frac{\sin x}{\cos x} = \tan x \end{aligned}$$

**✓ Progress Check**

Simplify the expression  $\frac{\csc \theta}{1 + \cot^2 \theta}$ .

**Answers**

$\sin \theta$

**EXAMPLE 2 VERIFYING AN IDENTITY**

Verify the identity

$$\sin \alpha - \sin^2 \alpha = \frac{1 - \sin \alpha}{\csc \alpha}$$

**SOLUTION**

$$\begin{aligned} \sin \alpha - \sin^2 \alpha &= \sin \alpha (1 - \sin \alpha) \\ &= \frac{1 - \sin \alpha}{\csc \alpha} \end{aligned}$$

**✓ Progress Check**

Verify the identity

$$\frac{\sin^2 y - 1}{1 - \sin y} = -1 - \sin y$$

**EXAMPLE 3 VERIFYING AN IDENTITY**

Verify the identity  $\cos x \tan x \csc x = 1$ .

**SOLUTION**

$$\begin{aligned} \cos x \tan x \csc x &= \cos x \left( \frac{\sin x}{\cos x} \right) \left( \frac{1}{\sin x} \right) \\ &= 1 \end{aligned}$$

✓ **Progress Check**

Verify the identity  $\sin x \sec x = \tan x$ .

**EXAMPLE 4** VERIFYING AN IDENTITY

Verify the identity

$$\frac{1}{1 - \sin x} + \frac{1}{1 + \sin x} = 2 \sec^2 x$$

**SOLUTION**

We begin with the left-hand side, combining fractions.

$$\begin{aligned} \frac{1}{1 - \sin x} + \frac{1}{1 + \sin x} &= \frac{1 + \sin x + 1 - \sin x}{(1 - \sin x)(1 + \sin x)} \\ &= \frac{2}{1 - \sin^2 x} = \frac{2}{\cos^2 x} \\ &= 2 \sec^2 x \end{aligned}$$

✓ **Progress Check**

Verify the identity  $\cos x + \tan x \sin x = \sec x$ .

**EXAMPLE 5** VERIFYING AN IDENTITY

Verify the identity

$$\frac{\cos \theta}{1 - \sin \theta} = \sec \theta + \tan \theta$$

**SOLUTION**

To obtain the expression  $1 - \sin^2 \theta$  in the denominator, we multiply numerator and denominator by  $1 + \sin \theta$ .

$$\begin{aligned} \frac{\cos \theta}{1 - \sin \theta} &= \left( \frac{\cos \theta}{1 - \sin \theta} \right) \left( \frac{1 + \sin \theta}{1 + \sin \theta} \right) \\ &= \frac{\cos \theta (1 + \sin \theta)}{1 - \sin^2 \theta} \\ &= \frac{\cos \theta (1 + \sin \theta)}{\cos^2 \theta} \\ &= \frac{1 + \sin \theta}{\cos \theta} \\ &= \frac{1}{\cos \theta} + \frac{\sin \theta}{\cos \theta} \\ &= \sec \theta + \tan \theta \end{aligned}$$

✓ **Progress Check**

Verify the identity

$$\frac{1 + \cos t}{\sin t} + \frac{\sin t}{1 + \cos t} = 2 \csc t$$

**EXAMPLE 6** VERIFYING AN IDENTITY

Verify the identity

$$\frac{\cot u - \tan u}{\sin u \cos u} = \csc^2 u - \sec^2 u$$

**SOLUTION**

We transform both sides of the equation by writing all trigonometric functions in terms of sine and cosine. For the left-hand side, we have

$$\begin{aligned} \frac{\cot u - \tan u}{\sin u \cos u} &= \frac{\frac{\cos u}{\sin u} - \frac{\sin u}{\cos u}}{\sin u \cos u} \\ &= \frac{\frac{\cos^2 u - \sin^2 u}{\sin u \cos u}}{\sin u \cos u} \end{aligned}$$

and for the right-hand side, we have

$$\begin{aligned} \csc^2 u - \sec^2 u &= \frac{1}{\sin^2 u} - \frac{1}{\cos^2 u} \\ &= \frac{\cos^2 u - \sin^2 u}{\sin^2 u \cos^2 u} \end{aligned}$$

We have successfully transformed both sides of the equation into the same expression. Since all the steps are reversible, we have verified the identity.

✓ **Progress Check**

Verify the identity

$$\frac{\sin x + \cos x}{\tan^2 x - 1} = \frac{\cos^2 x}{\sin x - \cos x}$$

**Exercise Set 8.1**

In Exercises 1–46, verify each of the identities.

- $\csc \gamma - \cos \gamma \cot \gamma = \sin \gamma$
- $\cot x \sec x = \csc x$

- $\sec v + \tan v = \frac{1 + \sin v}{\cos v}$
- $\cos \theta + \tan \theta \sin \theta = \sec \theta$
- $\sin \alpha \sec \alpha = \tan \alpha$

6.  $\sec \beta - \cos \beta = \sin \beta \tan \beta$
  7.  $3 - \sec^2 x = 2 - \tan^2 x$
  8.  $1 - 2 \sin^2 t = 2 \cos^2 t - 1$
  9.  $\frac{\sec^2 y}{\tan y} = \tan y + \cot y$
  10.  $\frac{\sin x + \cos x}{\cos x} = 1 + \tan x$
  11.  $\frac{\sin u}{\csc u} + \frac{\cos u}{\sec u} = 1$
  12.  $\frac{\tan^2 \alpha}{1 + \sec \alpha} = \sec \alpha - 1$
  13.  $\frac{\sec^2 \theta - 1}{\sec^2 \theta} = \sin^2 \theta$
  14.  $\sin^4 x + 2 \sin^2 x \cos^2 x + \cos^4 x = 1$
  15.  $\cos \gamma + \cos \gamma \tan^2 \gamma = \sec \gamma$
  16.  $\frac{1}{\tan u + \cot u} = \cos u \sin u$
  17.  $\frac{\sec w \sin w}{\tan w + \cot w} = \sin^2 w$
  18.  $(1 - \cos^2 \beta)(1 + \cot^2 \beta) = 1$
  19.  $(\sin \alpha + \cos \alpha)^2 + (\sin \alpha - \cos \alpha)^2 = 2$
  20.  $\frac{1 + \tan^2 u}{\csc^2 u} = \tan^2 u$
  21.  $\sec^2 v + \cos^2 v = \frac{\sec^4 v + 1}{\sec^2 v}$
  22.  $\sin^2 \theta - \tan^2 \theta = -\tan^2 \theta \sin^2 \theta$
  23.  $\frac{\sin^2 \alpha}{1 + \cos \alpha} = 1 - \cos \alpha$
  24.  $\cot x \sin^2 x = \cos x \sin x$
  25.  $\frac{\cos t}{1 + \sin t} = \frac{1 - \sin t}{\cos t}$
  26.  $\frac{\sin \beta}{1 + \cos \beta} + \frac{1 + \cos \beta}{\sin \beta} = 2 \csc \beta$
  27.  $\csc^2 \theta - \frac{\cos^2 \theta}{\sin^2 \theta} = 1$
  28.  $\frac{\cos^2 u}{1 - \sin u} = 1 + \sin u$
  29.  $\frac{\cot y}{1 + \cot^2 y} = \sin y \cos y$
  30.  $\frac{1 + \tan^2 x}{\tan^2 x} = \csc^2 x$
  31.  $\cos(-t) \csc(-t) = -\cot t$
  32.  $\sin(-\theta) \sec(-\theta) = -\tan \theta$
  33.  $\frac{\sec x + \csc x}{1 + \tan x} = \csc x$
  34.  $\frac{\sec u}{\sec u - 1} = \frac{1}{1 - \cos u}$
  35.  $\frac{1 + \tan x}{1 + \cot x} = \frac{\sec x}{\csc x}$
  36.  $(\tan u + \sec u)^2 = \frac{1 + \sin u}{1 - \sin u}$
  37.  $\frac{1 - \sin t}{1 + \sin t} = (\sec t - \tan t)^2$
  38.  $2 \csc^2 \theta - \csc^4 \theta = 1 - \cot^4 \theta$
  39.  $\frac{\sin^2 w}{\cot^4 w + \cos^2 w \sin^2 w} = \tan^2 w$
  40.  $\frac{\sin z + \tan z}{1 + \cos z} = \tan z$
  41.  $\frac{\sec \gamma - \csc \gamma}{\sec \gamma + \csc \gamma} = \frac{\tan \gamma - 1}{\tan \gamma + 1}$
  42.  $\frac{\cot x - 1}{1 - \tan x} = \frac{\csc x}{\sec x}$
  43.  $\frac{\tan \gamma - \sin \gamma}{\tan \gamma} = \frac{\sin^2 \gamma}{1 + \cos \gamma}$
  44.  $\cos^4 u - \sin^4 u = \cos^2 u - \sin^2 u$
  45.  $\frac{\csc x}{1 + \csc x} - \frac{\csc x}{1 - \csc x} = 2 \sec^2 x$
  46.  $\sin^3 \theta + \cos^3 \theta = (1 - \sin \theta \cos \theta)(\sin \theta + \cos \theta)$
- In Exercises 47–52, show that each of the equations is not an identity by finding a value of the variable for which the equation is not true.
47.  $\sin x = \sqrt{1 - \cos^2 x}$
  48.  $\tan x = \sqrt{\sec^2 x - 1}$
  49.  $(\sin t + \cos t)^2 = \sin^2 t + \cos^2 t$
  50.  $\sin \theta + \cos \theta = \sec \theta + \csc \theta$
  51.  $\sqrt{\cos^2 x} = \cos x$
  52.  $\sqrt{\cot^2 x} = \cot x$

## 8.2 The Addition and Subtraction Formulas

The identities that we verified in the examples and exercises of Section 8.1 were, in general, of no special significance. We were primarily interested in demonstrating manipulation with the fundamental identities. There are, however, many trigonometric identities that are indeed of importance. These identities are called **trigonometric formulas**.

Our objective in this section is to develop the **addition formulas** for  $\sin(s + t)$ ,  $\cos(s + t)$ , and  $\tan(s + t)$ , as well as the **subtraction formulas** for  $\sin(s - t)$ ,  $\cos(s - t)$ , and  $\tan(s - t)$ .


We begin with the derivation of  $\cos(s - t)$ . For convenience, assume that  $s$ ,  $t$ , and  $s - t$  are all positive and less than  $2\pi$ . Let  $P_1 = P(s)$ ,  $Q_1 = P(t)$ , and  $P_2 = P(s - t)$  be the points on the unit circle determined by  $s$ ,  $t$ , and  $s - t$ , respectively, as shown in Figure 1. Then  $\widehat{Q_2P_2Q_1P_1} = s$ ,  $\widehat{Q_2P_2Q_1} = t$ , and  $\widehat{Q_2P_2} = s - t$ . From Section 7.3, we know that the coordinates of  $P_1$ ,  $Q_1$ , and  $P_2$  are  $P_1 = P(\cos s, \sin s)$ ,  $Q_1 = P(\cos t, \sin t)$ , and  $P_2 = P(\cos(s - t), \sin(s - t))$ . Since the length of the arcs  $\widehat{Q_1P_1}$  and  $\widehat{Q_2P_2}$  are both equal (length of  $s - t$ ), the length of the chords  $\overline{Q_1P_1}$  and  $\overline{Q_2P_2}$  are also equal. By the distance formula, we have

$$\overline{Q_1P_1} = \overline{Q_2P_2}$$

$$\sqrt{(\cos s - \cos t)^2 + (\sin s - \sin t)^2} = \sqrt{[\cos(s - t) - 1]^2 + [\sin(s - t) - 0]^2}$$

Squaring both sides and rearranging terms, we have

$$\begin{aligned} \sin^2 s + \cos^2 s + \sin^2 t + \cos^2 t - 2 \cos s \cos t - 2 \sin s \sin t = \\ \sin^2(s - t) + \cos^2(s - t) - 2 \cos(s - t) + 1 \end{aligned}$$



**FIGURE 1** The Derivation of  $\cos(s - t)$



Since  $\sin^2 s + \cos^2 s = 1$ ,  $\sin^2 t + \cos^2 t = 1$  and  $\sin^2 (s - t) + \cos^2 (s - t) = 1$ , we have

$$2 - 2 \cos s \cos t - 2 \sin s \sin t = 2 - 2 \cos (s - t)$$

Solving for  $\cos (s - t)$  yields the formula

$$\cos (s - t) = \cos s \cos t + \sin s \sin t \quad (1)$$

### Graphing Calculator Power User's Corner 8.2



#### Verifying the Addition and Subtraction Formulas with the Calculator

The graphing calculator cannot take the place of algebraic manipulation in analytic trigonometry; however, it is a handy tool for testing potential values quickly. In this case we will examine  $\cos (s - t) = \cos s \cos t + \sin s \sin t$ . Let's assign  $s$  the value of  $\pi/3$  and let  $t = x$  for the sake of the calculator.

Press the  $Y=$  button and enter  $Y_1 = \cos(\pi/3 - x)$  and  $Y_2 = \cos(\pi/3)\cos(x) + \sin(\pi/3)\sin(x)$ . Graph these two equations in the trig viewing window. Repeat the procedure with  $s = \pi/4$ ,  $s = \pi/6$ , and  $s = 7\pi/6$ . What do you notice?

We obtain the formula for  $\cos (s + t)$  by writing

$$s + t = s - (-t)$$

Therefore

$$\begin{aligned} \cos (s + t) &= \cos (s - (-t)) \\ &= \cos s \cos (-t) + \sin s \sin (-t) \end{aligned}$$

Since  $\cos (-t) = \cos t$  and  $\sin (-t) = -\sin t$ ,

$$\cos (s + t) = \cos s \cos t - \sin s \sin t \quad (2)$$

### EXAMPLE 1 USING THE SUBTRACTION FORMULA

Find  $\cos 15^\circ$  without using tables or a calculator.

#### SOLUTION

Since  $15^\circ = 45^\circ - 30^\circ$ , we may use the formula for  $\cos (s - t)$  to obtain

$$\begin{aligned} \cos 15^\circ &= \cos (45^\circ - 30^\circ) \\ &= \cos 45^\circ \cos 30^\circ + \sin 45^\circ \sin 30^\circ \\ &= \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{3}}{2} + \frac{\sqrt{2}}{2} \cdot \frac{1}{2} \\ &= \frac{\sqrt{6} + \sqrt{2}}{4} \end{aligned}$$

✓ **Progress Check**

Solve Example 1 using  $15^\circ = 60^\circ - 45^\circ$ .

**EXAMPLE 2 USING THE ADDITION FORMULA**

Find the exact value of  $\cos \frac{5\pi}{12}$ .

**SOLUTION**

We note that  $\frac{5\pi}{12} = \frac{2\pi}{12} + \frac{3\pi}{12} = \frac{\pi}{6} + \frac{\pi}{4}$ . Then

$$\begin{aligned}\cos \frac{5\pi}{12} &= \cos \left( \frac{\pi}{6} + \frac{\pi}{4} \right) \\ &= \cos \frac{\pi}{6} \cos \frac{\pi}{4} - \sin \frac{\pi}{6} \sin \frac{\pi}{4} \\ &= \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2} - \frac{1}{2} \cdot \frac{\sqrt{2}}{2} \\ &= \frac{\sqrt{6} - \sqrt{2}}{4}\end{aligned}$$

✓ **Progress Check**

Solve Example 2 using the identity  $\frac{5\pi}{12} = \frac{9\pi}{12} - \frac{4\pi}{12}$ .

We may now establish the following relationships:

$$\cos \left( \frac{\pi}{2} - t \right) = \sin t \quad (3)$$

$$\sin \left( \frac{\pi}{2} - t \right) = \cos t \quad (4)$$

$$\tan \left( \frac{\pi}{2} - t \right) = \cot t \quad (5)$$

**Cofunctions**

As we stated in Section 7.2, Equations (6)–(11), sine and cosine are cofunctions, as are tangent and cotangent, and secant and cosecant. (Compare Equations (6), (7), and (8) of Section 7.2 with Equations (3), (4), and (5) above.)

Using the subtraction formula for cosine, we have

$$\begin{aligned}\cos \left( \frac{\pi}{2} - t \right) &= \cos \frac{\pi}{2} \cos t + \sin \frac{\pi}{2} \sin t \\ &= 0 \cdot \cos t + 1 \sin t \\ &= \sin t\end{aligned}$$

which establishes the identity in Equation (3). Replacing  $t$  with  $\frac{\pi}{2} - t$  in this identity yields

$$\begin{aligned}\cos\left[\frac{\pi}{2} - \left(\frac{\pi}{2} - t\right)\right] &= \sin\left(\frac{\pi}{2} - t\right) \\ \cos t &= \sin\left(\frac{\pi}{2} - t\right)\end{aligned}$$

which establishes the identity in Equation (4). The third identity follows from the definition of tangent and from Equations (3) and (4).

$$\tan\left(\frac{\pi}{2} - t\right) = \frac{\sin\left(\frac{\pi}{2} - t\right)}{\cos\left(\frac{\pi}{2} - t\right)} = \frac{\cos t}{\sin t} = \cot t$$

We will now prove the identity in Equation (6). (We leave the proof of Equation (7) as an exercise.)

$$\sin(s + t) = \sin s \cos t + \cos s \sin t \quad (6)$$

$$\sin(s - t) = \sin s \cos t - \cos s \sin t \quad (7)$$

From Equation (3),

$$\begin{aligned}\sin(s + t) &= \cos\left[\frac{\pi}{2} - (s + t)\right] \\ &= \cos\left[\left(\frac{\pi}{2} - s\right) - t\right] \\ &= \cos\left(\frac{\pi}{2} - s\right) \cos t + \sin\left(\frac{\pi}{2} - s\right) \sin t \\ &= \sin s \cos t + \cos s \sin t\end{aligned}$$

We conclude with the proof of the identity in Equation (8). (We leave the proof of Equation (9) as an exercise.)

$$\tan(s + t) = \frac{\tan s + \tan t}{1 - \tan s \tan t} \quad (8)$$

$$\tan(s - t) = \frac{\tan s - \tan t}{1 + \tan s \tan t} \quad (9)$$

$$\begin{aligned}
 \tan (s+t) &= \frac{\sin (s+t)}{\cos (s+t)} \\
 &= \frac{\sin s \cos t + \cos s \sin t}{\cos s \cos t - \sin s \sin t} \\
 &= \frac{\left(\frac{\sin s}{\cos s} \cdot \frac{\cos t}{\cos t}\right) + \left(\frac{\cos s}{\cos s} \cdot \frac{\sin t}{\cos t}\right)}{\left(\frac{\cos s}{\cos s} \cdot \frac{\cos t}{\cos t}\right) - \left(\frac{\sin s}{\cos s} \cdot \frac{\sin t}{\cos t}\right)} \\
 &= \frac{\tan s + \tan t}{1 - \tan s \tan t}
 \end{aligned}$$

### EXAMPLE 3 APPLYING THE ADDITION FORMULA

Show that  $\sin \left(x + \frac{3\pi}{2}\right) = -\cos x$ .

#### SOLUTION

$$\begin{aligned}
 \sin \left(x + \frac{3\pi}{2}\right) &= \sin x \cos \frac{3\pi}{2} + \cos x \sin \frac{3\pi}{2} \\
 &= (\sin x) 0 + (\cos x)(-1) \\
 &= -\cos x
 \end{aligned}$$

#### ✓ Progress Check

Verify that  $\tan (x - \pi) = \tan x$ .

### EXAMPLE 4 USING THE ADDITION FORMULA

Given  $\sin \alpha = -\frac{4}{5}$ , with  $\alpha$  an angle in quadrant III, and  $\cos \beta = -\frac{5}{13}$ , with  $\beta$  an angle in quadrant II, use the addition formula to find  $\sin (\alpha + \beta)$  and the quadrant in which  $\alpha + \beta$  lies.

#### SOLUTION

The addition formula

$$\sin (\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

requires that we know  $\sin \alpha$ ,  $\cos \alpha$ ,  $\sin \beta$ , and  $\cos \beta$ . Using the fundamental identity  $\sin^2 \alpha + \cos^2 \alpha = 1$ , we have

$$\cos^2 \alpha = 1 - \sin^2 \alpha = 1 - \frac{16}{25} = \frac{9}{25}$$

Taking the square root of both sides, we must have  $\cos \alpha = -\frac{3}{5}$  since  $\alpha$  is in quadrant III. Similarly,

$$\sin^2 \beta = 1 - \cos^2 \beta = 1 - \frac{25}{169} = \frac{144}{169}$$

Taking the square root of both sides, we must have  $\sin \beta = \frac{12}{13}$  since  $\beta$  is in quadrant II. Thus,

$$\begin{aligned}\sin(\alpha + \beta) &= \left(-\frac{4}{5}\right)\left(-\frac{5}{13}\right) + \left(-\frac{3}{5}\right)\left(\frac{12}{13}\right) \\ &= \frac{20}{65} - \frac{36}{65} = -\frac{16}{65}\end{aligned}$$

Since  $\sin(\alpha + \beta)$  is negative,  $\alpha + \beta$  lies in either quadrant III or quadrant IV. However, the sum of an angle that lies in quadrant III and an angle that lies in quadrant II cannot lie in quadrant III. Thus,  $\alpha + \beta$  lies in quadrant IV.

#### ✓ Progress Check

Given  $\cos \alpha = -\frac{4}{5}$ , with  $\alpha$  in quadrant III, and  $\cos \beta = \frac{3}{5}$ , with  $\beta$  in quadrant I, find  $\cos(\alpha - \beta)$  and the quadrant in which  $\alpha - \beta$  lies.

#### Answers

$-\frac{24}{25}$ , quadrant II

### Focus on Computing Sine and Cosine

```

10 LET S1 = 0.01745
20 LET C1 = 0.99985
30 PRINT "DEGREES",
   "SIN", "COS"
40 PRINT "1", S1, C1
50 LET S2 = S1
60 LET C2 = C1
70 FOR I = 2 TO 90
80 LET S3 = S2
90 LET S2 =
   (S1 * C2) +
   (C1 * S2)
100 LET C2 =
   (C1 * C2) -
   (S1 * S2)
110 PRINT I, S2, C2
120 NEXT I
130 END

```

We can make use of the trigonometric formulas to generate a table of sine and cosine values. Suppose we have determined that

$$\sin 1^\circ \approx 0.01745 \quad \cos 1^\circ \approx 0.99985 \quad (1)$$

We can then write

$$\sin(1^\circ + \alpha) = \sin 1^\circ \cos \alpha + \cos 1^\circ \sin \alpha$$

$$\cos(1^\circ + \alpha) = \cos 1^\circ \cos \alpha - \sin 1^\circ \sin \alpha$$

Substituting for  $\sin 1^\circ$  and  $\cos 1^\circ$  from (1),

$$\sin(1^\circ + \alpha) \approx 0.01745 \cos \alpha + 0.99985 \sin \alpha \quad (2)$$

$$\cos(1^\circ + \alpha) \approx 0.99985 \cos \alpha - 0.01745 \sin \alpha \quad (3)$$

Now, if we let  $\alpha = 1^\circ$ , (2) and (3) can be used to calculate  $\sin 2^\circ$  and  $\cos 2^\circ$ . We can then repeat the process with  $\alpha = 2^\circ$  to calculate  $\sin 3^\circ$  and  $\cos 3^\circ$ , and so on. We have provided a program in BASIC that will calculate sine and cosine values from  $2^\circ$  to  $90^\circ$  in increments of  $1^\circ$ .

## Exercise Set 8.2

In Exercises 1–6, show that the given equation is not an identity. (*Hint:* For each equation, find values of  $s$  and  $t$  for which the equation is not true.)

- $\cos(s - t) = \cos s - \cos t$
- $\sin(s + t) = \sin s + \sin t$
- $\sin(s - t) = \sin s - \sin t$
- $\cos(s + t) = \cos s + \cos t$
- $\tan(s + t) = \tan s + \tan t$
- $\tan(s - t) = \tan s - \tan t$

In Exercises 7–22, use the addition and subtraction formulas to find exact values.

- $\cos\left(\frac{\pi}{6} + \frac{\pi}{4}\right)$
- $\sin\left(\frac{\pi}{6} - \frac{\pi}{4}\right)$
- $\sin\left(\frac{\pi}{4} + \frac{\pi}{3}\right)$
- $\cos\left(\frac{\pi}{3} - \frac{\pi}{4}\right)$
- $\cos(30^\circ + 180^\circ)$
- $\tan(60^\circ + 300^\circ)$
- $\tan(300^\circ - 60^\circ)$
- $\sin(270^\circ - 45^\circ)$
- $\sin\frac{11\pi}{12}$  (*Hint:*  $\frac{11\pi}{12} = \frac{\pi}{6} + \frac{3\pi}{4}$ )
- $\tan\frac{7\pi}{12}$  (*Hint:*  $\frac{7\pi}{12} = \frac{\pi}{4} + \frac{\pi}{3}$ )
- $\cos\frac{7\pi}{12}$  (*Hint:*  $\frac{7\pi}{12} = \frac{5\pi}{6} - \frac{\pi}{4}$ )
- $\tan 75^\circ$  (*Hint:*  $75^\circ = 135^\circ - 60^\circ$ )
- $\sin\frac{7\pi}{6}$
- $\cos\frac{5\pi}{6}$
- $\tan 15^\circ$
- $\tan 165^\circ$

In Exercises 23–28, write the given expression in terms of cofunctions of complementary angles.

- $\sin 47^\circ$
- $\cos 78^\circ$
- $\tan\frac{\pi}{6}$
- $\tan 84^\circ$
- $\cos\frac{\pi}{3}$
- $\sin 72^\circ 30'$
- If  $\sin t = -\frac{3}{5}$ , with  $t$  in quadrant III, find  $\sin\left(\frac{\pi}{2} - t\right)$ .
- If  $\cos t = -\frac{5}{13}$ , with  $t$  in quadrant II, find  $\sin(t - \pi)$ .

- If  $\tan \theta = \frac{4}{3}$  and angle  $\theta$  lies in quadrant III, find  $\tan\left(\theta + \frac{\pi}{4}\right)$ .
- If  $\sec \theta = \frac{5}{3}$  and angle  $\theta$  lies in quadrant I, find  $\sin\left(\theta + \frac{\pi}{6}\right)$ .
- If  $\cos t = 0.4$ , with  $t$  in quadrant IV, find  $\tan(t + \pi)$ .
- If  $\sec \alpha = 1.2$  and angle  $\alpha$  lies in quadrant IV, find  $\tan(\alpha - \pi)$ .
- If  $\sin s = \frac{3}{5}$  and  $\cos t = -\frac{12}{13}$ , with  $s$  in quadrant II and  $t$  in quadrant III, find  $\sin(s + t)$ .
- If  $\sin s = -\frac{4}{5}$  and  $\csc t = \frac{13}{5}$ , with  $s$  in quadrant IV and  $t$  in quadrant II, find  $\cos(s - t)$ .
- If  $\cos \alpha = \frac{5}{13}$  and  $\tan \beta = -2$ , with angle  $\alpha$  in quadrant I and angle  $\beta$  in quadrant II, find  $\tan(\alpha + \beta)$ .
- If  $\sec \alpha = \frac{5}{3}$  and  $\cot \beta = \frac{15}{8}$ , with angle  $\alpha$  in quadrant IV and angle  $\beta$  in quadrant III, find  $\tan(\alpha - \beta)$ .

In Exercises 39–54, prove each of the following identities by transforming the left-hand side of the equation into the expression on the right-hand side.

- $\sin 2\alpha = 2 \sin \alpha \cos \alpha$
- $\cos 2t = \cos^2 t - \sin^2 t$
- $\tan 2\alpha = \frac{2 \tan \alpha}{1 - \tan^2 \alpha}$
- $\sin(x + y) \sin(x - y) = \sin^2 x - \sin^2 y$
- $\cos(x - y) \cos(x + y) = \cos^2 x \cos^2 y - \sin^2 x \sin^2 y$
- $\frac{\sin(s + t)}{\sin(s - t)} = \frac{\tan s + \tan t}{\tan s - \tan t}$
- $\csc\left(t + \frac{\pi}{2}\right) = \sec t$
- $\tan(\alpha + 90^\circ) = -\cot \alpha$
- $\tan\left(x + \frac{\pi}{4}\right) = \frac{1 + \tan x}{1 - \tan x}$
- $\csc(t - \pi) = -\csc t$
- $\cot(s - t) = \frac{1 + \tan s \tan t}{\tan s - \tan t}$
- $\cot(u + v) = \frac{\cot u \cot v - 1}{\cot u + \cot v}$

51.  $\sin(s + t) + \sin(s - t) = 2 \sin s \cos t$

52.  $\cos(s + t) + \cos(s - t) = 2 \cos s \cos t$

53. 
$$\frac{\sin(x + b) - \sin x}{b} = \sin x \left( \frac{\cos b - 1}{b} \right) + \cos x \left( \frac{\sin b}{b} \right)$$

54. 
$$\frac{\cos(x + b) - \cos x}{b} = \cos x \left( \frac{\cos b - 1}{b} \right) - \sin x \left( \frac{\sin b}{b} \right)$$

## 8.3 Double-Angle and Half-Angle Formulas

### Double-Angle Formulas

Our initial objective in this section is to derive expressions for  $\sin 2t$ ,  $\cos 2t$ , and  $\tan 2t$  in terms of trigonometric functions of  $t$ . We will establish the following double-angle formulas.

$$\sin 2t = 2 \sin t \cos t \quad (1)$$

$$\cos 2t = \cos^2 t - \sin^2 t \quad (2)$$

$$\tan 2t = \frac{2 \tan t}{1 - \tan^2 t} \quad (3)$$

To establish Equation (1), we rewrite  $2t$  as  $(t + t)$  and use the addition formula.

$$\begin{aligned} \sin 2t &= \sin(t + t) \\ &= \sin t \cos t + \cos t \sin t \\ &= 2 \sin t \cos t \end{aligned}$$

We proceed in the same manner to prove Equation (2).

$$\begin{aligned} \cos 2t &= \cos(t + t) \\ &= \cos t \cos t - \sin t \sin t \\ &= \cos^2 t - \sin^2 t \end{aligned}$$

Using the addition formula for the tangent function yields a proof of Equation (3).

$$\begin{aligned} \tan 2t &= \tan(t + t) \\ &= \frac{\tan t + \tan t}{1 - \tan t \tan t} \\ &= \frac{2 \tan t}{1 - \tan^2 t} \end{aligned}$$

#### EXAMPLE 1 USING THE DOUBLE-ANGLE FORMULAS

If  $\cos t = -\frac{3}{5}$  and  $P(t)$  is in quadrant II, evaluate  $\sin 2t$  and  $\cos 2t$ . In which quadrant does  $P(2t)$  lie?

**SOLUTION**

We first find  $\sin t$  by use of the fundamental identity  $\sin^2 t + \cos^2 t = 1$ . Thus,

$$\sin^2 t + \frac{9}{25} = 1$$

$$\sin^2 t = \frac{16}{25}$$

Since  $P(t)$  is in quadrant II,  $\sin t$  must be positive. Therefore,

$$\sin t = \frac{4}{5}$$

Applying the double-angle formulas with  $\cos t = -\frac{3}{5}$  and  $\sin t = \frac{4}{5}$ , we have

$$\sin 2t = 2 \sin t \cos t = 2\left(\frac{4}{5}\right)\left(-\frac{3}{5}\right) = -\frac{24}{25}$$

$$\cos 2t = \cos^2 t - \sin^2 t = \frac{9}{25} - \frac{16}{25} = -\frac{7}{25}$$

Since  $\sin 2t$  and  $\cos 2t$  are both negative, we conclude that  $P(2t)$  lies in quadrant III.

**✓ Progress Check**

If  $\sin \theta = \frac{5}{13}$  and  $\theta$  is in quadrant I, evaluate  $\sin 2\theta$  and  $\tan 2\theta$ .

**Answers**

$$\sin 2\theta = \frac{120}{169}, \tan 2\theta = \frac{120}{119}$$

**EXAMPLE 2 USING THE ADDITION AND DOUBLE-ANGLE FORMULAS**

Express  $\sin 3t$  in terms of  $\sin t$  and  $\cos t$ .

**SOLUTION**

We write  $3t$  as  $(2t + t)$ . Then

$$\begin{aligned} \sin 3t &= \sin (2t + t) \\ &= \sin 2t \cos t + \cos 2t \sin t \\ &= 2 \sin t \cos t \cos t + (\cos^2 t - \sin^2 t) \sin t \\ &= 2 \sin t \cos^2 t + \sin t \cos^2 t - \sin^3 t \\ &= 3 \sin t \cos^2 t - \sin^3 t \end{aligned}$$



**✓ Progress Check**Express  $\cos 3t$  in terms of  $\sin t$  and  $\cos t$ .**Answers**

$$\cos 3t = \cos^3 t - 3 \sin^2 t \cos t$$

If we begin with the formula for  $\cos 2t$  and use the fundamental identity  $\cos^2 t = 1 - \sin^2 t$ , we obtain

$$\begin{aligned}\cos 2t &= \cos^2 t - \sin^2 t \\ &= (1 - \sin^2 t) - \sin^2 t \\ &= 1 - 2 \sin^2 t\end{aligned}$$

Similarly, replacing  $\sin^2 t$  by  $1 - \cos^2 t$  yields

$$\begin{aligned}\cos 2t &= \cos^2 t - \sin^2 t \\ &= \cos^2 t - (1 - \cos^2 t) \\ &= 2 \cos^2 t - 1\end{aligned}$$

We then have two additional formulas for  $\cos 2t$ .

$$\cos 2t = 1 - 2 \sin^2 t \quad (4)$$

$$\cos 2t = 2 \cos^2 t - 1 \quad (5)$$

**EXAMPLE 3 DOUBLE-ANGLE FORMULAS IN VERIFYING AN IDENTITY**

Verify the identity

$$\frac{1 - \cos 2\alpha}{2 \sin \alpha \cos \alpha} = \tan \alpha$$

**SOLUTION**

Substituting  $\cos 2\alpha = 1 - 2 \sin^2 \alpha$ , we have

$$\begin{aligned}\frac{1 - \cos 2\alpha}{2 \sin \alpha \cos \alpha} &= \frac{1 - (1 - 2 \sin^2 \alpha)}{2 \sin \alpha \cos \alpha} \\ &= \frac{2 \sin^2 \alpha}{2 \sin \alpha \cos \alpha} \\ &= \frac{\sin \alpha}{\cos \alpha} \\ &= \tan \alpha\end{aligned}$$

✓ **Progress Check**

Verify the identity

$$\frac{1 + \cos 2\theta}{\sin 2\theta} = \cot \theta$$



**WARNING**

Note that

$$\frac{\sin 2t}{2} \neq \sin t$$

From Equation (1),

$$\frac{\sin 2t}{2} = \frac{2 \sin t \cos t}{2} = \sin t \cos t$$

## Half-Angle Formulas

If we begin with the alternative forms for  $\cos 2t$  given in Equations (4) and (5), we can obtain the following expressions for  $\sin^2 t$  and  $\cos^2 t$ . The expressions are often used in calculus.

$$\sin^2 t = \frac{1 - \cos 2t}{2} \quad (6)$$

$$\cos^2 t = \frac{1 + \cos 2t}{2} \quad (7)$$

We will use the identities in Equations (6) and (7) to derive formulas for  $\sin \frac{t}{2}$ ,  $\cos \frac{t}{2}$ , and  $\tan \frac{t}{2}$ . Substituting  $s = 2t$  into Equations (6) and (7), we obtain

$$\sin^2 \frac{s}{2} = \frac{1 - \cos s}{2}$$

$$\cos^2 \frac{s}{2} = \frac{1 + \cos s}{2}$$

Replacing  $s$  with  $t$  and solving, we have

$$\sin \frac{t}{2} = \pm \sqrt{\frac{1 - \cos t}{2}} \quad (8)$$

$$\cos \frac{t}{2} = \pm \sqrt{\frac{1 + \cos t}{2}} \quad (9)$$

The appropriate sign to use in Equations (8) and (9) depends on the quadrant in which  $P(\frac{t}{2})$  is located. Thus,  $\sin \frac{t}{2}$  is positive if  $P(\frac{t}{2})$  lies in quadrant I or II. Similarly, we choose the positive root for  $\cos \frac{t}{2}$  in Equation (9) if  $P(\frac{t}{2})$  lies in quadrant I or IV.

Using the identity

$$\tan \frac{t}{2} = \frac{\sin \frac{t}{2}}{\cos \frac{t}{2}}$$

we obtain

$$\tan \frac{t}{2} = \pm \sqrt{\frac{1 - \cos t}{1 + \cos t}} \quad (10)$$

Formulas (8), (9), and (10) are known as the **half-angle formulas**.

#### EXAMPLE 4 APPLYING THE HALF-ANGLE FORMULAS

Find the exact values of  $\sin 22.5^\circ$  and  $\cos 112.5^\circ$ .

#### SOLUTION

Applying the half-angle formulas with  $22.5^\circ = \frac{45^\circ}{2}$ , we have

$$\begin{aligned} \sin 22.5^\circ &= \sin \frac{45^\circ}{2} \\ &= \sqrt{\frac{1 - \cos 45^\circ}{2}} = \sqrt{\frac{1 - \frac{\sqrt{2}}{2}}{2}} \\ &= \frac{\sqrt{2 - \sqrt{2}}}{2} \end{aligned}$$

Note that we choose the positive square root since  $22.5^\circ$  is in the first quadrant and the sine function is positive in the first quadrant. Similarly,

$$\begin{aligned} \cos 112.5^\circ &= \cos \frac{225^\circ}{2} \\ &= -\sqrt{\frac{1 + \cos 225^\circ}{2}} \\ &= -\sqrt{\frac{1 - \cos 45^\circ}{2}} = -\sqrt{\frac{1 - \frac{\sqrt{2}}{2}}{2}} \\ &= -\frac{\sqrt{2 - \sqrt{2}}}{2} \end{aligned}$$

The negative square root was selected since  $112.5^\circ$  is in the second quadrant and the cosine function is negative in quadrant II.

✓ **Progress Check**

Use the half-angle formulas to evaluate  $\tan \frac{3\pi}{8}$ .

**Answers**

$$\sqrt{2} + 1$$

**EXAMPLE 5 APPLYING THE HALF-ANGLE FORMULAS**

If  $\sin \theta = -\frac{3}{5}$  and  $\theta$  is in quadrant III, evaluate  $\cos \frac{\theta}{2}$ .

**SOLUTION**

We first evaluate  $\cos \theta$  by using the identity

$$\cos^2 \theta = 1 - \sin^2 \theta = 1 - \frac{9}{25} = \frac{16}{25}$$

Since  $\theta$  is in quadrant III,  $\cos \theta$  is negative. Thus,  $\cos \theta = -\frac{4}{5}$ . We can now employ the half-angle formula

$$\begin{aligned} \cos \frac{\theta}{2} &= \pm \sqrt{\frac{1 + \cos \theta}{2}} \\ &= \pm \sqrt{\frac{1 - \frac{4}{5}}{2}} \\ &= \pm \frac{\sqrt{10}}{10} \end{aligned}$$

Since  $180^\circ < \theta < 270^\circ$ , we see that  $90^\circ < \frac{\theta}{2} < 135^\circ$ . Thus,  $\frac{\theta}{2}$  is in quadrant II and  $\cos \frac{\theta}{2}$  is negative. We conclude that

$$\cos \frac{\theta}{2} = -\frac{\sqrt{10}}{10}$$

✓ **Progress Check**

If  $\tan \alpha = \frac{3}{4}$  and  $\alpha$  is in quadrant III, evaluate  $\tan \frac{\alpha}{2}$ .

**Answers**

$$-3$$

### Exercise Set 8.3

In Exercises 1–12, use the given conditions to determine the value of the specified trigonometric function.

- If  $\sin u = \frac{3}{5}$  and  $P(u)$  is in quadrant II, find  $\cos 2u$ .
- If  $\cos x = -\frac{5}{13}$  and  $P(x)$  is in quadrant III, find  $\sin 2x$ .
- If  $\sec \alpha = -2$  and  $\alpha$  is in quadrant II, find  $\sin 2\alpha$ .
- If  $\tan \theta = \frac{4}{3}$  and  $\theta$  is in quadrant I, find  $\cos 2\theta$ .
- If  $\csc t = -\frac{17}{8}$  and  $P(t)$  is in quadrant IV, find  $\tan 2t$ .
- If  $\cot \beta = \frac{3}{4}$  and  $\beta$  is in quadrant III, find  $\cot 2\beta$ .
- If  $\sin 2\alpha = -\frac{4}{5}$  and  $2\alpha$  is in quadrant IV, find  $\sin 4\alpha$ .
- If  $\sec 5x = -\frac{13}{12}$  and  $P(5x)$  is in quadrant III, find  $\tan 10x$ .
- If  $\cos \frac{\theta}{2} = \frac{8}{17}$  and  $\frac{\theta}{2}$  is acute, find  $\cos \theta$ .
- If  $\csc \frac{t}{2} = -\frac{13}{5}$  and  $P(\frac{t}{4})$  is in quadrant IV, find  $\cos \frac{t}{2}$ .
- If  $\sin 42^\circ \approx 0.67$ , find  $\cos 84^\circ$ .
- If  $\cos 77^\circ \approx 0.22$ , find  $\cos 154^\circ$ .

In Exercises 13–18, use the half-angle formulas to find exact values for each of the following.

- |                          |                            |
|--------------------------|----------------------------|
| 13. $\sin 15^\circ$      | 14. $\cos 75^\circ$        |
| 15. $\tan \frac{\pi}{8}$ | 16. $\sec \frac{5\pi}{8}$  |
| 17. $\csc 165^\circ$     | 18. $\cot \frac{7\pi}{12}$ |

In Exercises 19–26, use the given conditions to determine the exact value of the specified trigonometric function.

- If  $\sin \theta = -\frac{4}{5}$  and  $\theta$  is in quadrant IV, find  $\cos \frac{\theta}{2}$ .

- If  $\cos \theta = \frac{3}{5}$  and  $\theta$  is in quadrant I, find  $\sin \frac{\theta}{2}$ .
- If  $\sec t = -3$  and  $P(t)$  is in quadrant II, find  $\sin \frac{t}{2}$ .
- If  $\tan x = \frac{4}{3}$  and  $P(x)$  is in quadrant III, find  $\cos \frac{x}{2}$ .
- If  $\cot \beta = \frac{3}{4}$  and  $\beta$  is in quadrant III, find  $\tan \frac{\beta}{2}$ .
- If  $\csc \alpha = \frac{13}{5}$  and  $\alpha$  is in quadrant II, find  $\tan \frac{\alpha}{2}$ .
- If  $\cos 4x = \frac{1}{3}$  and  $P(4x)$  is in quadrant IV, find  $\cos 2x$ .
- If  $\sec 6\alpha = -\frac{13}{12}$  and  $\alpha$  is in quadrant III, find  $\sin 3\alpha$ .

In Exercises 27–46, verify the identity.

- $\sin 50x = 2 \sin 25x \cos 25x$
- $(\sin \theta + \cos \theta)^2 = 1 + \sin 2\theta$
- $\tan 2y = \frac{2 \cot y}{\csc^2 y - 2}$
- $2 \sin^2 2t + \cos 4t = 1$
- $\sin 4\alpha = 4 \sin \alpha \cos^3 \alpha - 4 \sin^3 \alpha \cos \alpha$
- $\cos 4\beta = 1 - 8 \sin^2 \beta \cos^2 \beta$
- $\cos 2u = \frac{1 - \tan^2 u}{1 + \tan^2 u}$
- $\sin 2\theta = \frac{2 \tan \theta}{1 + \tan^2 \theta}$
- $\sin \frac{t}{2} \cos \frac{t}{2} = \frac{\sin t}{2}$
- $\tan \frac{y}{2} = \csc y - \cot y$
- $\sin \alpha - \cos \alpha \tan \frac{\alpha}{2} = \tan \frac{\alpha}{2}$
- $\frac{1 - \cos 2\beta}{1 + \cos 2\beta} = \tan^2 \beta$
- $\cos^4 x - \sin^4 x = \cos 2x$
- $\frac{\sin 2t}{\sin t} - \frac{\cos 2t}{\cos t} = \sec t$

41.  $\frac{2 \tan \alpha}{1 + \tan^2 \alpha} = \sin 2\alpha$

42.  $\cos^2 \frac{x}{2} = \frac{\tan x + \sin x}{2 \tan x}$

43.  $\sec 2t = \frac{\sec^2 t}{2 - \sec^2 t}$

44.  $\cos 2t + \cot 2t = \cot 2t (\sin t + \cos t)^2$

45.  $\tan \frac{t}{2} = \frac{1 - \cos t}{\sin t}$

46.  $\tan \frac{t}{2} = \frac{\sin t}{1 + \cos t}$

In Exercises 47–50, find the requested value.

47.  $\sin \left( 2 \arccos \frac{3}{5} \right)$

48.  $\cos \left( 2 \sin^{-1} \frac{3}{5} \right)$

49.  $\tan \left( 2 \arcsin \frac{5}{13} \right)$

50.  $\cos \left( 2 \arctan \frac{12}{5} \right)$

## 8.4 The Product-Sum Formulas

The objective of this section is to derive formulas that can transform sums of sines and cosines into products of sines and cosines, and vice versa. We use the word “sum” in a more general way to include the word “difference,” since subtraction can be thought of as adding a negative quantity.

The following formulas express a product as a sum.

$$\sin s \cos t = \frac{\sin (s+t) + \sin (s-t)}{2} \quad (1)$$

$$\cos s \sin t = \frac{\sin (s+t) - \sin (s-t)}{2} \quad (2)$$

$$\cos s \cos t = \frac{\cos (s+t) + \cos (s-t)}{2} \quad (3)$$

$$\sin s \sin t = \frac{\cos (s-t) - \cos (s+t)}{2} \quad (4)$$

### Product-Sum Formulas

To prove the identity in Equation (1), we begin with the right-hand side of the equation.

$$\begin{aligned} \frac{\sin (s+t) + \sin (s-t)}{2} &= \frac{(\sin s \cos t + \cos s \sin t) + (\sin s \cos t - \cos s \sin t)}{2} \\ &= \frac{2 \sin s \cos t}{2} \\ &= \sin s \cos t \end{aligned}$$

The proofs of the identities in Equations (2), (3), and (4) are very similar.

### Graphing Calculator Power User's Corner 8.4



#### Verifying the Product-Sum Formulas with the Calculator

As was mentioned in section 8.2, the graphing calculator cannot take the place of algebraic manipulation in analytic trigonometry. In this section, however, we will use it to look at  $\sin s \cos t$ . Let  $s = \pi/6$  and  $t = x$  for the purposes of the calculator and enter the left side of the formula into  $Y_1$  and the right hand side into  $Y_2$ . Be sure to enclose the numerator of the fraction in a set of parentheses. Graph these two equations in the trig viewing window and look for points of intersection. Repeat the procedure with various values of  $s$ . How could you extend this approach to the other product-sum formulas?

#### EXAMPLE 1 APPLYING THE PRODUCT-SUM FORMULAS

Express  $\sin 4x \cos 3x$  as a sum or a difference.

#### SOLUTION

Applying Equation (1), we obtain

$$\begin{aligned}\sin 4x \cos x &= \frac{\sin (4x + 3x) + \sin (4x - 3x)}{2} \\ &= \frac{\sin 7x + \sin x}{2}\end{aligned}$$

#### ✓ Progress Check

Express  $\sin 5x \sin 2x$  as a sum or as a difference.

#### Answers

$$\frac{1}{2}(\cos 3x - \cos 7x)$$

#### EXAMPLE 2 APPLYING THE PRODUCT-SUM FORMULAS

Evaluate the product  $\cos \frac{5\pi}{8} \cos \frac{3\pi}{8}$  by a product-sum formula.

#### SOLUTION

Using Equation (3), we have

$$\begin{aligned}\cos \frac{5\pi}{8} \cos \frac{3\pi}{8} &= \frac{1}{2} \left[ \cos \left( \frac{5\pi}{8} + \frac{3\pi}{8} \right) + \cos \left( \frac{5\pi}{8} - \frac{3\pi}{8} \right) \right] \\ &= \frac{1}{2} \left[ \cos \pi + \cos \frac{\pi}{4} \right] \\ &= \frac{1}{2} \left[ -1 + \frac{\sqrt{2}}{2} \right] = \frac{\sqrt{2} - 2}{4}\end{aligned}$$

✓ **Progress Check**

Evaluate  $\cos \frac{\pi}{3} \sin \frac{\pi}{6}$  by a product-sum formula.

**Answers**

$$\frac{1}{4}$$

The following formulas express a sum as a product.

$$\sin s + \sin t = 2 \sin \frac{s+t}{2} \cos \frac{s-t}{2} \quad (5)$$

$$\sin s - \sin t = 2 \cos \frac{s+t}{2} \sin \frac{s-t}{2} \quad (6)$$

$$\cos s + \cos t = 2 \cos \frac{s+t}{2} \cos \frac{s-t}{2} \quad (7)$$

$$\cos s - \cos t = -2 \sin \frac{s+t}{2} \sin \frac{s-t}{2} \quad (8)$$

To prove the identity in Equation (5), we begin with the right-hand side and apply Equation (1). Then

$$\begin{aligned} 2 \sin \frac{s+t}{2} \cos \frac{s-t}{2} &= 2 \left\{ \frac{1}{2} \left[ \sin \left( \frac{s+t}{2} + \frac{s-t}{2} \right) + \sin \left( \frac{s+t}{2} - \frac{s-t}{2} \right) \right] \right\} \\ &= \sin s + \sin t \end{aligned}$$

This establishes Equation (5). The other identities are established in a similar fashion.

**EXAMPLE 3 APPLYING THE PRODUCT-SUM FORMULAS**

Express  $\sin 5x - \sin 3x$  as a product.

**SOLUTION**

Using Equation (6), we have

$$\begin{aligned} \sin 5x - \sin 3x &= 2 \cos \frac{5x+3x}{2} \sin \frac{5x-3x}{2} \\ &= 2 \cos 4x \sin x \end{aligned}$$

✓ **Progress Check**

Express  $\cos 6x + \cos 2x$  as a product.

**Answers**

$$2 \cos 4x \cos 2x$$



**EXAMPLE 4** APPLYING THE PRODUCT-SUM FORMULASEvaluate  $\cos \frac{5\pi}{12} - \cos \frac{\pi}{12}$  using a product-sum-formula.**SOLUTION**

Using Equation (8), we have

$$\begin{aligned}\cos \frac{5\pi}{12} - \cos \frac{\pi}{12} &= -2 \sin \frac{\pi}{4} \sin \frac{\pi}{6} \\ &= -2 \left( \frac{\sqrt{2}}{2} \right) \frac{1}{2} = -\frac{\sqrt{2}}{2}\end{aligned}$$

**✓ Progress Check**

Evaluate

$$\sin \frac{11\pi}{12} - \sin \frac{5\pi}{12}$$

using a product-sum formula.

**Answers**

$$-\frac{\sqrt{2}}{2}$$

**Exercise Set 8.4**

In Exercises 1–8, express each product as a sum or difference.

1.  $2 \sin 5\alpha \cos \alpha$
2.  $-3 \cos 6x \sin 2x$
3.  $\sin 3x \sin (-2x)$
4.  $\cos 7t \cos (-3t)$
5.  $-2 \cos 2\theta \cos 5\theta$
6.  $\sin \frac{5\theta}{2} \sin \frac{\theta}{2}$
7.  $\cos (\alpha + \beta) \cos (\alpha - \beta)$
8.  $-\sin 2u \cos 4u$

In Exercises 9–12, evaluate each product by using a product-sum formula.

9.  $\cos \frac{7\pi}{8} \sin \frac{5\pi}{8}$
10.  $\cos \frac{\pi}{3} \cos \frac{\pi}{6}$
11.  $\sin 120^\circ \cos 60^\circ$
12.  $\sin \frac{13\pi}{12} \sin \frac{11\pi}{12}$

In Exercises 13–20, express each sum or difference as a product.

13.  $\sin 5x + \sin x$
14.  $\cos 8t - \cos 2t$

15.  $\cos 2\theta + \cos 6\theta$

16.  $\sin 5\alpha - \sin 7\alpha$

17.  $\sin (\alpha + \beta) + \sin (\alpha - \beta)$

18.  $\cos \frac{x}{2} - \cos \frac{3x}{2}$

19.  $\sin 7x - \sin 3x$

20.  $\cos 5\theta + \cos 3\theta$

In Exercises 21–24, evaluate each sum by using a product-sum formula.

21.  $\cos 75^\circ + \cos 15^\circ$

22.  $\sin \frac{5\pi}{12} + \sin \frac{\pi}{12}$

23.  $\cos \frac{3\pi}{4} - \cos \frac{\pi}{4}$

24.  $\sin \frac{13\pi}{12} - \sin \frac{5\pi}{12}$

In Exercises 25–34, verify the identities.

25.  $\sin 40^\circ + \sin 20^\circ = \cos 10^\circ$

26.  $\cos 70^\circ - \cos 10^\circ = -\sin 40^\circ$

27.  $\frac{\sin 5\theta - \sin 3\theta}{\cos 3\theta - \cos 5\theta} = \cot 4\theta$

28.  $\frac{\cos 7x - \cos x}{\sin 7x + \sin x} = -\tan 3x$

$$29. \frac{\sin t - \sin s}{\cos t - \cos s} = -\cot \frac{s+t}{2}$$

$$30. \frac{\sin s + \sin t}{\cos s + \cos t} = \tan \frac{s+t}{2}$$

$$31. \frac{\sin 50^\circ - \sin 10^\circ}{\cos 50^\circ - \cos 10^\circ} = -\sqrt{3}$$

$$32. 2 \sin \left(\theta + \frac{\pi}{4}\right) \sin \left(\theta - \frac{\pi}{4}\right) = -\cos 2\theta$$

$$33. \frac{\cot x - \tan x}{\cot x + \tan x} = \cos 2x$$

$$34. \cos 6x \cos 2x + \sin^2 4x = \cos^2 2x$$

35. Express  $\sin ax \cos bx$  as a sum.

36. Express  $\cos ax \cos bx$  as a sum.

37. Prove the product-sum formulas given in Equations (2), (3), and (4).

38. Prove the product-sum formulas given in Equations (6), (7), and (8).

## 8.5 Trigonometric Equations

Thus far, this chapter has dealt exclusively with trigonometric identities. We now seek to solve trigonometric equations that may be true for some values of the variable but not for all values.

We have seen that algebraic equations may have just one or two solutions. The situation is quite different with trigonometric equations. Since trigonometric functions are periodic by nature, if we find one solution, there must be an infinite number of solutions. To deal with this situation, we first seek all solutions  $t$  such that  $0 \leq t < 2\pi$ . Then, for every integer  $n$ ,  $t + 2\pi n$  is also a solution. The following example illustrates this procedure for finding the solution set.

### EXAMPLE 1 SOLVING A TRIGONOMETRIC EQUATION

Find all solutions of the equation  $\cos t = 0$ .

#### SOLUTION

The only values in the interval  $[0, 2\pi)$  for which  $\cos t = 0$  are  $\frac{\pi}{2}$  and  $\frac{3\pi}{2}$ . Then every solution is included among those values of  $t$  such that

$$t = \frac{\pi}{2} + 2\pi n \quad \text{or} \quad t = \frac{3\pi}{2} + 2\pi n, \quad n \text{ an integer}$$

Since  $\frac{3\pi}{2} = \frac{\pi}{2} + \pi$ , the solution set can be written in a more compact form as

$$t = \frac{\pi}{2} + \pi n, \quad n \text{ an integer}$$

Factoring provides an important technique for solving trigonometric equations. If we can write the equation in the form  $P(x)Q(x) = 0$ , we can find the solutions by setting  $P(x) = 0$  and  $Q(x) = 0$ .

It may also be helpful to think of substituting a new variable for some trigonometric expression. Thus, the equation

$$4 \sin^2 x + 3 \sin x - 1 = 0$$

can be viewed as a quadratic in  $u$

$$4u^2 + 3u - 1 = 0$$

by substituting  $u = \sin x$ .

### Graphing Calculator Power User's Corner 8.5



#### Solving Trigonometric Equations

Take a simple equation, such as  $\tan \theta = \frac{1}{2}$ . The calculator can help you solve this easily with a little manipulation. Go to  $Y=$  and enter  $Y_1 = \tan(x)$  and  $Y_2 = \frac{1}{2}$ . Now graph in the standard trig window and look for points of intersection of the two functions. Generally a problem will ask for all solutions within a certain set of  $x$  values such as  $[-\pi, \pi]$ . What are the solutions to this equation?

The same approach can be used for more difficult equations, such as  $\sin 2\theta - 3 \sin \theta = 0$ . In this case it would be beneficial to view the equation as  $\sin 2\theta = 3 \sin \theta$  and enter the left side in  $Y_1$  and the right side in  $Y_2$ . Once again look for the points of intersection.

### EXAMPLE 2 RESTRICTING THE SOLUTIONS TO A TRIGONOMETRIC EQUATION

Find all solutions of the equation  $2 \cos^2 t - \cos t - 1 = 0$  in the interval  $[0, 2\pi)$ .

#### SOLUTION

Factoring the left side of the equation yields

$$(2 \cos t + 1)(\cos t - 1) = 0$$

Setting each factor equal to 0, we have

$$2 \cos t + 1 = 0 \quad \text{or} \quad \cos t - 1 = 0$$

so that

$$\cos t = -\frac{1}{2} \quad \text{or} \quad \cos t = 1$$

The solutions of  $\cos t = -\frac{1}{2}$  in the interval  $[0, 2\pi)$  are  $t = \frac{2\pi}{3}$  and  $t = \frac{4\pi}{3}$ . The only solution of  $\cos t = 1$  in the interval  $[0, 2\pi)$  is  $t = 0$ . Thus, the solutions of the original problem are

$$t = \frac{2\pi}{3}, \quad t = \frac{4\pi}{3}, \quad \text{and} \quad t = 0$$

### ✓ Progress Check

Find all solutions of the equation  $2 \sin^2 t - 3 \sin t + 1 = 0$  in the interval  $[0, 2\pi)$ .

### Answers

$$\frac{\pi}{6}, \frac{5\pi}{6}, \frac{\pi}{2}$$

If the solutions of the trigonometric equation are angles, the answer may be given in either radians or degrees.

### EXAMPLE 3 EXPRESSING SOLUTIONS IN RADIANS AND DEGREES

Find all solutions of the equation  $\tan \theta \cos^2 \theta - \tan \theta = 0$ .

### SOLUTION

Factoring the left side yields

$$(\tan \theta)(\cos^2 \theta - 1) = 0$$

Setting each factor equal to 0,

$$\tan \theta = 0 \quad \text{or} \quad \cos^2 \theta = 1$$

so that

$$\tan \theta = 0, \quad \cos \theta = 1, \quad \text{or} \quad \cos \theta = -1$$

These equations yield the following solutions in the interval  $[0, 2\pi)$ .

$$\tan \theta = 0 : \quad \theta = 0 \quad \text{or} \quad \theta = \pi$$

$$\cos \theta = 1 : \quad \theta = 0$$

$$\cos \theta = -1 : \quad \theta = \pi$$

The solutions of the original equation are

$$\theta = 0 + 2\pi n \quad \text{and} \quad \theta = \pi + 2\pi n, \quad n \text{ an integer}$$

which can be expressed more compactly as

$$\theta = \pi n, \quad n \text{ an integer}$$

In degree measure, the solutions are

$$\theta = 180^\circ n, \quad n \text{ an integer}$$

**EXAMPLE 4 EXPRESSING RESTRICTED SOLUTIONS IN RADIANS AND DEGREES**

Find all solutions of the equation  $\sin 2\theta - 3 \sin \theta = 0$  in the interval  $[0, 2\pi)$ .

**SOLUTION**

Using the identity  $\sin 2\theta = 2 \sin \theta \cos \theta$ , we have

$$2 \sin \theta \cos \theta - 3 \sin \theta = 0$$

$$\sin \theta (2 \cos \theta - 3) = 0$$

$$\sin \theta = 0 \quad \text{or} \quad 2 \cos \theta - 3 = 0$$

$$\sin \theta = 0 \quad \text{or} \quad \cos \theta = \frac{3}{2}$$

The equation  $\cos \theta = \frac{3}{2}$  has no solutions. The solutions of  $\sin \theta = 0$  are  $\theta = 0$  and  $\theta = \pi$ . Therefore, the solutions of the original equation are

$$\theta = 0 \quad \text{and} \quad \theta = \pi$$

or, in degree measure,

$$\theta = 0^\circ \quad \text{and} \quad \theta = 180^\circ$$

**✓ Progress Check**

Find all solutions of the equation  $\cos 2\theta + \cos \theta = 0$ .

**Answers**

$$\frac{\pi}{3} + 2\pi n, \pi + 2\pi n, \frac{5\pi}{3} + 2\pi n \quad \text{or}$$

$$60^\circ + 360^\circ n, 180^\circ + 360^\circ n, 300^\circ + 360^\circ n$$

Equations involving multiple angles can often be solved by using a substitution of variable. The following example shows what may occur when seeking solutions in the interval  $[0, 2\pi)$ .

**EXAMPLE 5 SUBSTITUTION OF A VARIABLE IN TRIGONOMETRIC EQUATIONS**

Find all solutions of the equation  $\cos 3x = 0$  in the interval  $[0, 2\pi)$ .

**SOLUTION**

We are given

$$\cos 3x = 0, \quad 0 \leq x < 2\pi$$

Substituting  $t = 3x$ , we obtain

$$\cos t = 0, \quad 0 \leq \frac{t}{3} < 2\pi$$

or

$$\cos t = 0, \quad 0 \leq t < 6\pi$$

Note that we seek solutions of  $\cos t = 0$  in the interval  $[0, 6\pi)$  rather than  $[0, 2\pi)$ . The solutions are then

$$t = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \frac{7\pi}{2}, \frac{9\pi}{2}, \frac{11\pi}{2}$$

Since  $x = \frac{t}{3}$ , we obtain

$$x = \frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{3\pi}{2}, \frac{11\pi}{6}$$



### WARNING

When you perform a substitution of variable, you must remember to go back and express the answers in terms of the original variable.

### EXAMPLE 6 SUBSTITUTION OF VARIABLE

Find all solutions of the equation

$$3 \tan^2 x + \tan x - 1 = 0$$

in the interval  $[0, \pi)$ .

### SOLUTION

Since the equation does not factor easily, consider  $u = \tan x$ . Then we obtain

$$3u^2 + u - 1 = 0$$

From the quadratic formula,

$$u = \frac{-1 \pm \sqrt{13}}{6}$$

so that

$$\tan x = \frac{-1 \pm \sqrt{13}}{6}$$

Therefore,

$$\tan x \approx 0.4342586 \quad \text{and} \quad \tan x \approx -0.7675919$$

in which case

$$x \approx \tan^{-1} 0.4342586 \approx 0.4096865$$

and

$$x \approx \tan^{-1} (-0.7675919) \approx -0.6546652$$

Although the first value for  $x$  is in the interval  $[0, \pi)$ , the second value for  $x$  is not. In fact, this second value is in the interval  $(-\frac{\pi}{2}, \frac{\pi}{2})$ , the range of  $\tan^{-1} x$ . Since the period of  $\tan x$  is  $\pi$ ,

$$\begin{aligned} x &\approx -0.6546652 + \pi \\ &\approx 2.4869275 \end{aligned}$$

is also a solution. Observe that this new value for  $x$  is in the interval  $[0, \pi)$ .

### Graphing Calculator Power User's Corner 8.5b



#### Solving Trigonometric Equations Graphically


The ZOOM-IN capability of your graphing calculator enables you to solve trigonometric equations that cannot be solved algebraically.

#### EXAMPLE


Find all solutions of the equation  $\sin x = x^2 - 1$ .

#### SOLUTION


A graph in the viewing rectangle  $-2\pi \leq X \leq 2\pi$  and  $-3 \leq Y \leq 3$  indicates that there are two solutions to the equation  $\sin x = x^2 - 1$ , as shown in Figure 2(a). TRACEing to the points of intersection approximates the solutions as  $X_1 \approx -0.6$  and  $X_2 \approx 1.4$ . To obtain greater accuracy, use the ZOOM BOX method, as shown in Figures 2(b) and 2(c).



(a)  $-6.28319 \leq X \leq 6.28319$   
 $-3 \leq Y \leq 3$   
 $XSCL = 1, YSCL = 1$



(b)  $-0.63674 \leq X \leq -0.63672$   
 $-0.59458 \leq Y \leq -0.59456$   
 $XSCL = 1, YSCL = 1$   
 $X_1 \approx -0.6367$



(c)  $1.40962 \leq X \leq 1.40963$   
 $0.98703 \leq Y \leq 0.98705$   
 $XSCL = 1, YSCL = 1$   
 $X_2 \approx 1.4096$

FIGURE 2 Graphs of  $y = \sin x$  and  $y = x^2 - 1$

### Exercise Set 8.5

In Exercises 1–20, find all solutions of the given equation in the interval  $[0, 2\pi)$ . Express the answers in both radian measure and degree measure.

1.  $2 \sin \theta - 1 = 0$
2.  $2 \cos \theta + 1 = 0$
3.  $\cos \alpha + 1 = 0$
4.  $\cot \gamma + 1 = 0$
5.  $4 \cos^2 \alpha = 3$
6.  $\tan^2 \theta = 3$
7.  $3 \tan^2 \alpha = 1$
8.  $2 \cos^2 \alpha - 1 = 0$
9.  $2 \sin^2 \beta = \sin \beta$
10.  $\sin \alpha = \cos \alpha$
11.  $2 \cos^2 \theta - 3 \cos \theta + 1 = 0$
12.  $2 \sin^2 \theta - \sin \theta - 1 = 0$
13.  $\sin 5\theta = 1$
14.  $\tan 3\beta = -\sqrt{3}$
15.  $2 \sin^2 \alpha - 3 \cos \alpha = 0$
16.  $\csc 2\theta = 2$
17.  $2 \cos^2 \theta - 1 = \sin \theta$
18.  $\cos^2 2\alpha = \frac{1}{4}$
19.  $\sin^2 \beta + 3 \cos \beta - 3 = 0$
20.  $2 \cos^2 \theta \tan \theta - \tan \theta = 0$

In Exercises 21–38, find all the solutions of the given equation.

21.  $3 \tan^2 x - 1 = 0$
22.  $2 \sin^2 y - 1 = 0$
23.  $3 \cot^2 \theta - 1 = 0$
24.  $1 - 4 \cos^2 t = 0$
25.  $\sec 2u - 2 = 0$
26.  $\tan 3x - 1 = 0$
27.  $\sin 4x = 0$
28.  $\cos 5t = -1$

29.  $4 \cos^2 2t - 3 = 0$
30.  $\csc^2 2x - 2 = 0$
31.  $\sin 2t + 2 \cos t = 0$
32.  $\sin 2t + 3 \cos t = 0$
33.  $\cos 2t + \sin t = 0$
34.  $2 \cos 2t + 2 \sin t = 0$
35.  $\tan^2 x - \tan x = 0$
36.  $\sec^2 x - 3 \sec x + 2 = 0$
37.  $2 \sin^2 x + 3 \sin x - 2 = 0$
38.  $2 \cos^2 x - 5 \cos x - 3 = 0$



In Exercises 39–42, find the approximate solutions of the given equations in the interval  $[0, 2\pi)$  by using a calculator.

39.  $5 \sin^2 x - \sin x - 2 = 0$
40.  $\sec^2 y - 5 \sec y + 6 = 0$
41.  $3 \tan^2 u + 5 \tan u + 1 = 0$
42.  $\cos^2 t - 2 \sin t + 3 = 0$



In Exercises 43–46, find the approximate solutions of the given equation in the interval  $[0, 2\pi)$  by ZOOMING-IN on the point(s) of intersection of appropriate graphs on your graphing calculator.

43.  $\cos x = x$
44.  $\sin x = \cos x$
45.  $\tan x = 8 - \frac{1}{2}x^2$
46.  $3 \cos \frac{x}{2} = x^2 - 3$





# Chapter Summary

## Terms and Symbols

addition formulas	536	half-angle formulas	547	trigonometric equations	529
cofunctions	539	identities	529	trigonometric expressions	529
double-angle formulas	543	product-sum formulas	551	trigonometric formulas	536
fundamental identities	530	subtraction formulas	536	trigonometric identities	529

## Key Ideas for Review

Topic	Page	Key Idea
<b>Trigonometric Identity</b>	<b>529</b>	A trigonometric identity is true for all values that may be assumed by the variable.
<i>Fundamental Identities</i>	530	The fundamental identities are trigonometric identities that are directly related to the definitions of the trigonometric functions.
<i>Verification of Identities</i>	532	The fundamental identities can be used to verify other trigonometric identities. The techniques frequently used to verify identities include: <ol style="list-style-type: none"> <li>1. factoring</li> <li>2. writing trigonometric functions in terms of sine and cosine</li> <li>3. performing some of the indicated operations and simplifying complicated expressions</li> <li>4. multiplying numerator and denominator of some fraction involving trigonometric functions by some trigonometric expression to obtain forms such as <math>1 - \sin^2 \theta</math>, <math>1 - \cos^2 \theta</math> or <math>\sec^2 \theta - 1</math>, which can be further simplified.</li> </ol>

**Trigonometric Formulas** 536 Some of the most useful trigonometric formulas are the following.

### Addition Formulas

$$\sin(s + t) = \sin s \cos t + \cos s \sin t$$

$$\cos(s + t) = \cos s \cos t - \sin s \sin t$$

$$\tan(s + t) = \frac{\tan s + \tan t}{1 - \tan s \tan t}$$

### Double-Angle Formulas

$$\sin 2t = 2 \sin t \cos t$$

$$\cos 2t = \cos^2 t - \sin^2 t$$

$$\tan 2t = \frac{2 \tan t}{1 - \tan^2 t}$$

## Key Ideas for Review

Topic	Page	Key Idea
<b>Half-Angle Formulas</b>		
		$\sin \frac{t}{2} = \pm \sqrt{\frac{1 - \cos t}{2}}$ $\cos \frac{t}{2} = \pm \sqrt{\frac{1 + \cos t}{2}}$ $\tan \frac{t}{2} = \pm \sqrt{\frac{1 - \cos t}{1 + \cos t}}$
<b>Trigonometric Equations</b>	555	Since the trigonometric functions are periodic, a trigonometric equation has either no solution or an infinite number of solutions.
<i>Restricted Solutions</i>	556	Sometimes the solutions to a trigonometric equation are restricted to a specific interval.

## Review Exercises

Solutions to exercises whose numbers are in **blue** are in the Solutions section in the back of the book.

In Exercises 1–3, verify the given identity.

1.  $\sin \theta \sec \theta + \tan \theta = 2 \tan \theta$

2.  $\frac{\cos^2 x}{1 - \sin x} = 1 + \sin x$

3.  $\sin \alpha + \sin \alpha \cot^2 \alpha = \csc \alpha$

In Exercises 4–7, determine the exact value of the given expression by using the addition formulas.

4.  $\sin \left( \frac{\pi}{6} + \frac{\pi}{4} \right)$

5.  $\cos (45^\circ + 90^\circ)$

6.  $\tan \left( \frac{\pi}{3} + \frac{\pi}{4} \right)$

7.  $\sin \frac{7\pi}{12}$

In Exercises 8–11, write the given expression in terms of cofunctions of complementary angles.

8.  $\csc 15^\circ$

9.  $\cos 23^\circ$

10.  $\sin \frac{\pi}{8}$

11.  $\tan \frac{2\pi}{7}$

12. If  $\cos \theta = -\frac{12}{13}$  and  $0 \leq \theta \leq 180^\circ$  find  $\sin (\pi - \theta)$ .

13. If  $\sec \alpha = \frac{5}{4}$  and  $\alpha$  lies in quadrant IV, find  $\csc \left( \alpha + \frac{\pi}{3} \right)$ .

14. If  $\sin t = -\frac{3}{5}$  and  $P(t)$  is in quadrant III, find  $\tan (t + \pi)$ .

15. If  $\cos \alpha = -\frac{12}{13}$  and  $\tan \beta = -\frac{5}{2}$ , with angles  $\alpha$  and  $\beta$  in quadrant II, find  $\tan (\alpha + \beta)$ .

16. If  $\sin x = \frac{3}{5}$  and  $\csc y = \frac{13}{12}$ , with  $P(x)$  in quadrant II and  $P(y)$  in quadrant I, find  $\cos (x - y)$ .

17. If  $\csc u = -\frac{5}{4}$  and  $P(u)$  is in quadrant IV, find  $\cos 2u$ .

18. If  $\tan \alpha = -\frac{3}{4}$  and  $0 \leq \alpha \leq 180^\circ$  find  $\sin 2\alpha$ .

19. If  $\sin 2t = \frac{3}{5}$  and  $P(2t)$  is in quadrant I, find  $\sin 4t$ .

20. If  $\sin \theta = 0.5$  and  $\frac{\pi}{2} \leq \theta \leq \pi$  find  $\sin 2\theta$ .

21. If  $\cos \frac{\theta}{2} = \frac{12}{13}$  and  $\theta$  is acute, find  $\sin \theta$ .

22. If  $\sin \alpha = -\frac{3}{5}$  and  $\alpha$  is in quadrant III, find  $\cos \frac{\alpha}{2}$ .

## Review Exercises

23. If  $\cot t = -\frac{4}{3}$  and  $P(t)$  is in quadrant IV, find  $\tan \frac{t}{2}$ .
24. If  $\cos 4x = \frac{2}{3}$  and  $P(4x)$  is in quadrant IV, find  $\cos 2x$ .
25. Find the exact value of  $\cos 15^\circ$  by using a half-angle formula.
26. Find the exact value of  $\sin \frac{\pi}{8}$  by using a half-angle formula.
27. Find the exact value of  $\tan 112.5^\circ$  by using a half-angle formula.

In Exercises 28–30, verify the given identity.

28.  $\cos 30x = 1 - 2 \sin^2 15x$
29.  $\frac{1}{2} \sin 2y = \frac{\sin y}{\sec y}$       30.  $\tan \frac{\alpha}{2} = \frac{1 - \cos \alpha}{\sin \alpha}$
31. Express  $\sin \frac{3\alpha}{2} \sin \frac{\alpha}{2}$  as a sum or difference.
32. Express  $\cos 3x - \cos x$  as a product.
33. Evaluate  $\sin 75^\circ \sin 15^\circ$  by using a product-sum formula.

34. Evaluate  $\cos \frac{3\pi}{4} + \cos \frac{\pi}{4}$  by using a product-sum formula.

In Exercises 35–37, find all solutions of the given equation in the interval  $[0, 2\pi)$ . Express the answers in radian measure.

35.  $2 \cos^2 \alpha - 1 = 0$
36.  $2 \sin \theta \cos \theta = 0$
37.  $\sin 2t - \sin t = 0$

In Exercises 38–40, find all solutions of the given equation. Express the answers in degree measure.

38.  $\cos^2 \alpha - 2 \cos \alpha = 0$
39.  $\tan 3x + 1 = 0$
40.  $4 \sin^2 2t = 3$



In 41. Find the approximate solutions of  $x \sin x = 10 - x^2$  in the interval  $[0, 2\pi)$  by ZOOMING-IN on the points of intersection of appropriate graphs on your graphing calculator.

## Review Test

1. Verify the identity  $4 - \tan^2 x = 5 - \sec^2 x$ .

In Exercises 2 and 3, determine exact values of the given expressions by using the addition formulas.

2.  $\cos (270^\circ + 30^\circ)$       3.  $\tan \left( \frac{\pi}{4} - \frac{\pi}{3} \right)$
4. Write  $\sin 47^\circ$  in terms of its cofunction.
5. If  $\cos \theta = \frac{4}{5}$  and  $\theta$  lies in quadrant IV, find  $\sin (\theta - \pi)$ .
6. If  $\sin x = -\frac{5}{13}$  and  $\tan y = \frac{8}{3}$  with angles  $x$  and  $y$  in quadrant III, find  $\tan (x - y)$ .
7. If  $\sin v = -\frac{12}{13}$  and  $P(v)$  is in quadrant IV, find  $\cos 2v$ .
8. If  $\cos 2\alpha = -\frac{4}{5}$  and  $2\alpha$  is in quadrant II, find  $\cos 4\alpha$ .

9. If  $\csc \alpha = -2$  and  $\alpha$  is in quadrant III, find  $\cos \frac{\alpha}{2}$ .
10. Find the exact value of  $\tan 15^\circ$  by using a half-angle formula.
11. Verify the identity
- $$\sin \frac{x}{4} = 2 \sin \frac{x}{8} \cos \frac{x}{8}$$
12. Express  $\sin 2x + \sin 3x$  as a product.
13. Express  $\sin 150^\circ - \sin 30^\circ$  by using a product-sum formula.
14. Find all solutions of the equation  $4 \sin^2 \alpha = 3$  in the interval  $[0, 2\pi)$ . Express the answers in radian measure.
15. Find all solutions of the equation  $\sin^2 \theta - \cos^2 \theta = 0$  and express the answers in degree measure.

